arXiv:math-ph/0703040v1 13 Mar 2007

Solution of the Radial Schrödinger Equation for the Potential Family $V(r)=\frac{A}{r^2}-\frac{B}{r}+Cr^{\kappa}$ using the Asymptotic Iteration Method

M. Aygun†, O. Bayrak‡§ and I. Boztosun§

†*Faculty of Arts and Sciences, Department of Physics,*
Ataturk University, Erzurum, Turkey
‡*Faculty of Arts and Sciences, Department of Physics,*
Bozok University, Yozgat, Turkey and
§*Faculty of Arts and Sciences, Department of Physics,*
Erciyes University, Kayseri, Turkey

(Dated: February 3, 2008)

Abstract

We present the exact and iterative solutions of the radial Schrödinger equation for a class of potential, $V(r)=\frac{A}{r^2}-\frac{B}{r}+Cr^{\kappa}$, for various values of κ from -2 to 2, for any n and l quantum states by applying the asymptotic iteration method. The global analysis of this potential family by using the asymptotic iteration method results in exact analytical solutions for the values of $\kappa=0,-1$ and -2. Nevertheless, there are no analytical solutions for the cases $\kappa=1$ and 2. Therefore, the energy eigenvalues are obtained numerically. Our results are in excellent agreement with the previous works.

PACS numbers: 03.65.Ge

Keywords: asymptotic iteration method, eigenvalues and eigenfunctions, Kratzer, Modified Kratzer, Goldman-Krivchenkov, spiked harmonic oscillator, Coulomb plus linear and Coulomb plus harmonic oscillator potentials.

I. INTRODUCTION

We search for the exact solution of a new class of potential as in the following form:

$$V(r) = \frac{A}{r^2} - \frac{B}{r} + Cr^{\kappa} \tag{1}$$

This potential includes various potentials according to the values of the potential parameters:

a) By choosing $C = 0$, this potential turns into the Kratzer potential, which is analytically solvable by using various methods [1, 2, 3]. This potential has been used extensively so far in order to describe the molecular structure and interactions [1, 2, 3].

b) The Gol'dman-Krivchenkov potential is obtained by choosing $A \neq 0$, $B = 0$ and $\kappa = 2$ and it is solved analytically by using various methods [2, 4]. Furthermore, this potential becomes the spiked harmonic oscillator potential [4] when $A = 0$, $B = 1$ and $\kappa = 2$.

c) The Coulomb plus linear potential form is obtained by choosing $A = 0$ and $\kappa = 1$ and is solved by methods such as the envelope and variational methods [4]. This potential has raised great interest in atomic and molecular physics (Ref. [5] and references therein).

d) The Coulomb plus harmonic oscillator potential form is obtained by choosing $A = 0$ and $\kappa = 2$. This form is solved by using the moment method [6], the shifted $1/N$ expansion method [7] and the envelope method [4]. In addition, it is applied to examine the Zeeman quadratic effect [8] and the magnetic field effect in the hydrogen atom [9].

These potentials have been extensively used to describe the bound and the continuum states of the interactions systems and a great number of papers have been published for the exact and numerical solutions of these potentials (see Refs. [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13] and references therein). Thus, it would be interesting and important to solve the non-relativistic radial Schrödinger equation for a new solvable potential family for any n and l quantum states. Recently, an alternative method, called as the Asymptotic Iteration Method (AIM), has been developed by Çiftçi *et al.* [13] for solving the second-order homogeneous linear differential equations and it has been applied to solve the non-relativistic radial Schrödinger equation as well as the relativistic wave equations [13, 14, 15, 16, 17, 18]. In this paper, we aim to show that AIM could give the energy eigenvalues for the potentials that have analytical solutions obtained by using different methods. Moreover, AIM could provide energy eigenvalues for potentials that have no analytical solutions for any n and l quantum states with various κ values.

In the next section, we introduce AIM and then in section III, the analytical solution of the Schrödinger equation is obtained by using AIM for the potentials with $\kappa = 0, -1$ and -2 and a closed form for the energy eigenvalues and corresponding eigenfunctions for any n and l quantum numbers are given. Then, in section IV, the energy eigenvalues for a new solvable potential with $\kappa = 1$ and 2 cases by using AIM iteration procedure are obtained. Finally, in section V, we remark on these results.

II. OVERVIEW OF THE ASYMPTOTIC ITERATION METHOD (AIM)

AIM is proposed to solve the second-order differential equations of the form [13, 16].

$$y'' = \lambda_0(x)y' + s_0(x)y \tag{2}$$

where $\lambda_0(x) \neq 0$. The variables, $s_0(x)$ and $\lambda_0(x)$, are sufficiently differentiable. The differential equation (2) has a general solution [13]

$$y(x) = exp\left(-\int^{x} \alpha(x^{'})dx^{'}\right)\left[C_2 + C_1\int^{x} exp\left(\int^{x^{'}}[\lambda_0(x^{''}) + 2\alpha(x^{''})]dx^{''}\right)dx^{'}\right] \tag{3}$$

if $k > 0$, for sufficiently large k, we obtain the $\alpha(x)$ values from

$$\frac{s_k(x)}{\lambda_k(x)} = \frac{s_{k-1}(x)}{\lambda_{k-1}(x)} = \alpha(x), \quad k = 1, 2, 3, \ldots \tag{4}$$

where

$$\begin{aligned} \lambda_k(x) &= \lambda'_{k-1}(x) + s_{k-1}(x) + \lambda_0(x)\lambda_{k-1}(x) \\ s_k(x) &= s'_{k-1}(x) + s_0(x)\lambda_{k-1}(x), \qquad\qquad k = 1, 2, 3, \ldots \end{aligned} \tag{5}$$

It should be noted that one can also start the recurrence relations from $k = 0$ with the initial conditions $\lambda_{-1} = 1$ and $s_{-1} = 0$ [15]. For a given potential, the radial Schrödinger equation is converted to the form of equation (2). Then, $\mathrm{s}_0(x)$ and $\lambda_0(x)$ are determined and $\mathrm{s}_k(x)$ and $\lambda_k(x)$ parameters are calculated by the recurrence relations given by equation (5).

The energy eigenvalues are obtained from the roots of the quantization condition, given by the termination condition of the method in equation (4). The quantization condition of the method together with equation (5) can also be written as follows

$$\delta_k(x) = \lambda_k(x)s_{k-1}(x) - \lambda_{k-1}(x)s_k(x) = 0 \qquad k = 1, 2, 3, \ldots \tag{6}$$

The energy eigenvalues are obtained from this equation if the problem is exactly solvable. If not, for a specific n principal quantum number, we choose a suitable x_0 point, determined generally as the maximum value of the asymptotic wave function or the minimum value of the potential, and the approximate energy eigenvalues are obtained from the roots of this equation for sufficiently great values of k with iteration.

The wave functions are determined by using the following wave function generator

$$y_n(x) = C_2 exp(-\int^x \frac{s_k(x')}{\lambda_k(x')} dx') \tag{7}$$

where $k \geq n$, n represents the radial quantum number and k shows the iteration number. For exactly solvable potentials, the radial quantum number n is equal to the iteration number k and the eigenfunctions are obtained directly from equation (7). For nontrivial potentials that have no exact solutions, k is always greater than n in these numerical solutions and the approximate energy eigenvalues are obtained from the roots of equation (6) for sufficiently great values of k by iteration.

III. $\kappa = 0, -1$ AND -2 CASES: ANALYTICAL SOLUTIONS

Inserting the potential given by equation (1) into the Schrödinger equation gives

$$\frac{d^2R_{nl}}{dr^2} + \frac{2m}{\hbar^2}\left[E - \frac{A}{r^2} + \frac{B}{r} - Cr^{\kappa} - \frac{l(l+1)\hbar^2}{2mr^2}\right] R_{nl} = 0 \tag{8}$$

where n and l are radial and orbital angular momentum quantum numbers, A, B and C are strictly positive constants. By using the following *ansatze:*

$$-\varepsilon^2 = \frac{2mE}{\hbar^2}, \quad -\epsilon^2_{nl} = -\varepsilon^2_{nl} - \widetilde{C}, \quad \widetilde{A} = \frac{2mA}{\hbar^2}, \quad \widetilde{B} = \frac{2mB}{\hbar^2}, \quad \widetilde{C} = \frac{2mC}{\hbar^2}, \tag{9}$$

Equation (8) becomes

$$\frac{d^2R_{nl}}{dr^2} + \left[-\varepsilon^2 - \frac{\widetilde{A}}{r^2} + \frac{\widetilde{B}}{r} - \widetilde{C}r^{\kappa} - \frac{l(l+1)}{r^2}\right] R_{nl} = 0 \tag{10}$$

The aim of this paper is to show how to obtain the analytical and numerical solutions of $V(r) = \frac{A}{r^2} - \frac{B}{r} + Cr^{\kappa}$ potential with different κ values. κ=0, -1 and -2 values of this potential have analytical solutions and have been extensively studied so far [1, 2, 3, 18]. Therefore, we do not give the details of the calculations and only show the analytical solutions. By inserting $\kappa = 0, -1$ and -2 into equation (10) and by doing some simple algebra described in the previous section, the energy eigenvalues and regular eigenfunctions are obtained as follows:

κ	Eigenvalues	Eigenfunctions
0,	$E_{nl} = C - \frac{mB^2}{2\hbar^2}\left(n+\frac{1}{2}+\sqrt{(l+\frac{1}{2})^2+\frac{2mA}{\hbar^2}}\right)^{-2}$,	$R_{nl}(r) = Ne^{-\epsilon_{nl}^{\kappa=0}r}\ {}_1F_1(-n, 2\Lambda^{\kappa=0}+2; 2\epsilon_{nl}^{\kappa=0}r)$
-1,	$E_{nl} = -\frac{m(B-C)^2}{2\hbar^2}\left(n+\frac{1}{2}+\sqrt{(l+\frac{1}{2})^2+\frac{2mA}{\hbar^2}}\right)^{-2}$,	$R_{nl}(r) = Ne^{-\varepsilon_{nl}^{\kappa=-1}r}\ {}_1F_1(-n, 2\Lambda^{\kappa=-1}+2; 2\varepsilon_{nl}^{\kappa=-1}r)$
-2,	$E_{nl} = -\frac{mB^2}{2\hbar^2}\left(n+\frac{1}{2}+\sqrt{(l+\frac{1}{2})^2+\frac{2m}{\hbar^2}(A+C)}\right)^{-2}$,	$R_{nl}(r) = Ne^{-\varepsilon_{nl}^{\kappa=-2}r}\ {}_1F_1(-n, 2\Lambda^{\kappa=-2}+2; 2\varepsilon_{nl}^{\kappa=-2}r)$

with

$$\begin{array}{ll} \epsilon_{nl}^{\kappa=0} = \frac{\widetilde{B}}{2(n+\Lambda+1)}, & \Lambda^{\kappa=0} = -\frac{1}{2}+\sqrt{(l+\frac{1}{2})^2+\widetilde{A}} \\ \varepsilon_{nl}^{\kappa=-1} = \frac{\widetilde{B}-\widetilde{C}}{2(n+\Lambda+1)}, & \Lambda^{\kappa=-1} = -\frac{1}{2}+\sqrt{(l+\frac{1}{2})^2+\widetilde{A}} \\ \varepsilon_{nl}^{\kappa=-2} = \frac{\widetilde{B}}{2(n+\Lambda+1)}, & \Lambda^{\kappa=-2} = -\frac{1}{2}+\sqrt{(l+\frac{1}{2})^2+\widetilde{A}+\widetilde{C}} \end{array} \tag{11}$$

These results are in excellent agreement with the previous results obtained by using different methods (see [1, 2, 3, 18] and references therein).

IV. $\kappa = 1$ AND 2 CASES: ITERATIVE SOLUTIONS

For cases $\kappa = 1$ and 2, the exact analytical solutions can not be found and in this section, we present how to find the energy eigenvalues by applying the asymptotic iteration method. If we consider $V(r) = \frac{A}{r^2} - \frac{B}{r} + Cr^\kappa$ potential in the three-dimensional radial Schrödinger equation, we obtain equation (8). The straightforward application of AIM to solve this equation gives us the energy eigenvalues, however, we have observed that the energy eigenvalues oscillate and do not converge within a reasonable number of iteration. The sequence appears to converge when the number of iterations $k \simeq 30$, but then it begins to oscillate as the iteration number k increases. This result violates the principle behind the AIM; as the number of iteration increases, the method should converge and should not oscillate. In order to overcome this problem and obtain a rapid convergence, we make a change of variables as $r = r_0\rho$, where $r_0 = \frac{\hbar^2}{2mB}$, then we obtain

$$\frac{d^2R_{nl}}{d\rho^2} + \left[\varepsilon + \frac{1}{\rho} - \gamma^2\rho^\kappa - \frac{l'(l'+1)}{\rho^2}\right]R_{nl} = 0 \tag{12}$$

with the following *ansatz*

$$\varepsilon = \frac{\hbar^2 E}{2mB^2}, \quad \gamma^2 = \frac{2mCr_0^{\kappa+2}}{\hbar^2}, \quad l' = -\frac{1}{2} + \sqrt{(l+\frac{1}{2})^2 + \widetilde{A}}, \quad \widetilde{A} = \frac{2mA}{\hbar^2} \tag{13}$$

we can transform equation (12) to another Schrödinger equation form by changing the variable to $\rho = u^2$ and then by inserting $R(u) = u^{\frac{1}{2}}\phi(u)$ into the transformed equation. Therefore, we obtain the Schrödinger equation as follows

$$\frac{d^2\phi(u)}{du^2} + \left[4\varepsilon u^2 + 4 - 4\gamma^2 u^{2\kappa+2} - \frac{\Lambda(\Lambda+1)}{u^2}\right]\phi(u) = 0 \tag{14}$$

where $\Lambda = 2l' + \frac{1}{2}$.

A. $\kappa = 1$ Case

For $\kappa = 1$, we obtain the following equation by using equation (14):

$$\frac{d^2\phi(u)}{du^2} + \left[4\varepsilon u^2 + 4 - 4\gamma^2 u^4 - \frac{\Lambda(\Lambda+1)}{u^2}\right]\phi(u) = 0 \tag{15}$$

If we take the wave function in the following form

$$\phi(u) = u^{\Lambda+1} exp(-\frac{\gamma\beta u^4}{2}) f(u) \tag{16}$$

where β is an arbitrarily introduced constant to improve the convergence speed of the method [15]. We also introduce γ in the asymptotic wave function since the parameter γ appears explicitly in the potential. It is known that when changing the value of γ in the potential, it affects the eigenvalues and the shape of the wave function. Therefore, by introducing γ into equation (16), we control the change of the potential parameters and, as a result, obtain a better convergence. If we insert this wave function into equation (15), then we obtain the second-order homogeneous linear differential equation as follows;

$$\frac{d^2 f(u)}{du^2} = \left[2(2\beta\gamma u^3 - \frac{\Lambda+1}{u})\right]\frac{df(u)}{du} + \left[(4\beta\gamma\Lambda + 10\beta\gamma - 4\varepsilon)u^2 - 4\beta^2\gamma^2 u^6 - 4 + 4\gamma^2 u^4\right] f(u) \tag{17}$$

which is now amenable to an AIM solution. By comparing this equation with equation (2), we can write the $s_0(u)$ and $\lambda_0(u)$ values as below

$$\begin{aligned} s_0(u) &= (4\beta\gamma\Lambda + 10\beta\gamma - 4\varepsilon)u^2 - 4\beta^2\gamma^2 u^6 - 4 + 4\gamma^2 u^4 && (18) \\ \lambda_0(u) &= 2(2\beta\gamma u^3 - \frac{\Lambda+1}{u}) && (19) \end{aligned}$$

In order to obtain the energy eigenvalues from equation (17) by using equation (5), we obtain the $s_k(r)$ and $\lambda_k(r)$ in terms of $s_0(r)$ and $\lambda_0(r)$. Then, by using the quantization condition of the method given by equation (4), we obtain the energy eigenvalues. Therefore, we have to choose a suitable u_0 point to solve the equation $\delta_n(u_0, \varepsilon) = 0$ iteratively in order to find ε values. In this study, we obtain the u_0 from the maximum point of the asymptotic wave function, which is the same as the root of $\lambda_0(u) = 0$, thus $u_0 = \left(\frac{\Lambda+1}{2\beta\gamma}\right)^{1/4}$. This straightforward application of AIM gives us the energy eigenvalues as the sequence appears to converge when the number of iterations (k) are $k \simeq 30$ as shown in Table I.

In this table, we also present the convergence rate of AIM calculations. The energy eigenvalues appear as the number of iterations are $k \simeq 30$. However, the speed of the convergence depends on the arbitrarily introduced constant, β. We have investigated the optimum values of β that give the best convergence and have kept the one that appears to yield the best convergence rate. Therefore, for case κ=1, we have performed calculations for the different values of β. It is seen from Table I that the best convergence is obtained when the constant β values are β=0.4, 0.5, and 0.6. For other values of β, the convergence needs more iteration.

Having determined the value of β in an empirical way, for case κ=1, the energy eigenvalues by using AIM are shown for different values of n and l for convergence constant $\beta = 0.5$, $\widetilde{A} = 1$ and $\gamma = 1$ in Table II.

B. $\kappa = 2$ Case

For $\kappa = 2$, we obtain the following equation by using equation (14):

$$\frac{d^2\phi(u)}{du^2} + \left[4\varepsilon u^2 + 4 - 4\gamma^2 u^6 - \frac{\Lambda(\Lambda+1)}{u^2}\right]\phi(u) = 0 \tag{20}$$

If we take the wave function in the following form

$$\phi(u) = u^{\Lambda+1} exp(-\frac{\gamma\beta u^4}{2}) f(u) \tag{21}$$

When we insert this wave function into equation (20), we obtain the second-order homogeneous linear differential equation as follows

$$\frac{d^2 f(u)}{du^2} = 2\left[(2\beta\gamma u^3 - \frac{\Lambda+1}{u})\right]\frac{df(u)}{du} + \left[(4\beta\gamma\Lambda + 10\beta\gamma - 4\varepsilon)u^2 - 4\beta^2\gamma^2 u^6 - 4 + 4\gamma^2 u^6\right] f(u) \tag{22}$$

which is now amenable to an AIM solution. By comparing this equation with equation (2), we obtain $s_0(u)$ and $\lambda_0(u)$ values as below

$$s_0 = 4\beta\gamma u^2\Lambda - 4\beta^2\gamma^2 u^6 + 10\beta\gamma u^2 - 4\varepsilon u^2 - 4 + 4\gamma^2 u^6 \tag{23}$$

$$\lambda_0 = 2(2\beta\gamma u^3 - \frac{\Lambda + 1}{u}) \tag{24}$$

Similar to case $\kappa = 1$, in order to obtain the energy eigenvalues from equation (22) by using equation (5), we obtain the $s_k(r)$ and $\lambda_k(r)$ in terms of $s_0(r)$ and $\lambda_0(r)$. Then, by using the quantization condition of the method given by equation (4), we obtain the energy eigenvalues. For this case, we choose $u = (\frac{\Lambda+1}{2\beta\gamma})^{\frac{1}{4}}$ and we obtain the energy eigenvalues after the $k = 30$ iterations. The AIM results are presented in Table III for different values of n and l quantum numbers for different values of γ. In Table IV and V, we compare our results with previous works conducted for special forms of our potential. Our results are in excellent agreement with moment method [6], which is the closest one to exact solution. They are better than perturbation results.

V. CONCLUSION

In this paper, we have presented the exact and iterative solutions of the radial Schrödinger equation for a class of potential, $V(r) = \frac{A}{r^2} - \frac{B}{r} + Cr^\kappa$, for various values of κ from -2 to 2 for any n and l quantum states by applying the asymptotic iteration method. According to the value of κ, this potential family includes Kratzer, Modified Kratzer, Goldman-Krivchenkov or spiked harmonic oscillator, Coulomb plus linear and Coulomb plus harmonic oscillator potentials.

These potentials have been extensively used to describe the bound and the continuum states of the interactions systems and a great number of papers have been published for the exact and numerical solutions of these potentials (see Refs. [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13] and references therein). We have examined these potentials in this paper and have attempted to obtain exact or numerical solutions.

The global analysis of this potential family by using the asymptotic iteration method has resulted in exact analytical solutions for the values of $\kappa = 0, -1$ and -2. In addition, closed-forms for the energy eigenvalues as well as the corresponding eigenfunctions are obtained, but for $\kappa = 1$ and 2, there are no analytical solutions. Therefore, the energy eigenvalues are obtained numerically.

The advantage of the asymptotic iteration method is that it gives the eigenvalues directly by transforming the second-order differential equation into a form of $y'' = \lambda_0(r)y' + s_0(r)y$.

The wave functions are easily constructed by iterating the values of $s_0(r)$ and $\lambda_0(r)$. The asymptotic iteration method results in exact analytical solutions if there is and provides the closed-forms for the energy eigenvalues as well as the corresponding eigenfunctions. Where there is no such a solution, the energy eigenvalues are obtained by using an iterative approach [14, 15, 16, 17, 18, 19]. As it is presented, AIM puts no constraint on the potential parameter values involved and it is easy to implement. The results are sufficiently accurate for practical purposes. It is worth extending this method to examine other interacting systems.

Acknowledgments

This work is supported by the Scientific and Technical Research Council of Turkey (TÜBİTAK) under the project number TBAG-2398 and Erciyes University (FBT-04-16).

VI. REFERENCES

[1] A. Kratzer, Z. Phys. **3** (1920) 289.

[2] S. Flügge, `Practical Quantum Mechanics` Vol. I, Springer, Berlin, (1994).

[3] E. Kasap, B. Gonul and M. Simsek, Chem. Phys. Lett. **172** (1990) 499.

[4] H. Ciftci, R. L. Hall and Q. D. Katatbeh, J. Phys. A: Math. Gen. **36** (2003) 7001.

[5] E. Castro and P. Martin, J. Phys. A: Math. Gen. **33** (2000) 5321.

[6] D. Bessis, E. R. Vrscay and C. R. Handy, J. Phys. A: Math. Gen. **20** (1987) 419.

[7] R. K. Roychoudhury and Y. P. Varshni, J. Phys. A: Math. Gen. **21** (1988) 3025.

[8] J. E. Avron, Ann. Phys. **131** (1991) 73.

[9] M. Taut, J. Phys. A: Math. Gen. **28** (1995) 2081.

[10] J. Pliva, J. Mol. Spectrosc. **193** (1999) 193.

[11] P. G. Hajigeorgiou, J. Mol. Spectrosc. **235** (2006) 111.

[12] C. A. Singh and O. B. Devi, Int. J. Quantum Chem. **106** (2006) 415.

[13] H. Ciftci, R. L. Hall and N. Saad, J. Phys. A: Math. Gen. **36**(2003) 11807.

[14] T. Barakat, J. Phys. A: Math. Gen. **39** (2006) 823.

[15] F. M. Fernandez, J. Phys. A: Math. Gen. **37** (2004) 6173.

[16] O. Bayrak and I. Boztosun, J. Phys. A: Math. Gen. 39 (2006) 6955.

[17] H. Ciftci, R. L. Hall and N. Saad, Phys. Lett. **A340** (2005) 388.

[18] O. Bayrak and I. Boztosun, Int. J. Quantum Chem. (2006), in press.

[19] I. Boztosun, M. Karakoc, F. Yasuk and A. Durmus, Journal of Mathematical Physics **47** (2006) 062301.

k	$\beta = 0.2$	$\beta = 0.4$	$\beta = 0.5$	$\beta = 0.6$	$\beta = 0.7$	$\beta = 0.9$	$\beta = 2.0$
20	2.36068441	2.36071158	2.36071387	2.36071387	2.36080271	2.36168203	2.46417111
30	2.36071440	2.36071239	2.36071239	2.36071239	2.36071435	2.36076626	2.39133769
40	2.36071234	”	”	”	2.36071245	2.36071612	2.37021799
50	2.36071282	”	”	”	2.36071239	2.36071269	2.36372451
60	2.36071253	”	”	”	2.36071239	2.36071241	2.36168245
70	2.36071268	”	”	”	2.36071239	2.36071239	2.36102992
80	2.36071240	”	”	”	2.36071238	2.36071239	2.36081631
90	2.36071238	”	”	”	2.36071236	2.36071418	2.36076466

TABLE I: For $\kappa = 1$ case, the energy eigenvalues (ε) for the $n = 0$ and $l = 0$ states by means of several β values. We take $\widetilde{A} = 1$ and $\gamma = 1$.

k	$n = 0, l = 0$	$n = 1, l = 0$	$n = 1, l = 1$	$n = 2, l = 0$	$n = 2, l = 1$	$n = 2, l = 2$
20	2.36071387	4.112474	4.7245997	5.57024	6.097373	6.739515
30	2.36071239	4.112295	4.7242771	5.55292	6.074553	6.706985
40	2.36071239	4.112291	4.7242690	5.55211	6.073379	6.704987
50	”	4.112290	4.7242688	5.55208	6.073327	6.704887
60	”	”	”	5.55207	6.073324	6.704883
70	”	”	”	”	”	”

TABLE II: For $\kappa = 1$ case, the energy eigenvalues (ε) for the several quantum states. We take $\widetilde{A} = 1$, $\gamma = 1$ and convergence constant $\beta = 0.5$.

n	l	$\gamma = 0.1$	$\gamma = 1$	$\gamma = 10$
0	0	0.1220043681	3.3582483393	39.6495973187
	1	0.3258602332	4.8963878137	53.8428825862
	2	0.5473302077	6.7964025443	72.0047051977
1	0	0.5720226793	7.4731840675	79.9800946486
	1	0.7518135075	8.9639368105	94.0443338946
	2	0.9621813797	10.8379872414	112.1314195487
2	0	0.9982832867	11.5427585197	120.1879082158
	1	1.1684333950	13.0102255287	134.1850120876
	2	1.372953445	14.8691668817	152.2273430525

TABLE III: For $\kappa = 2$ case, the energy eigenvalues (ε) for several quantum numbers. We take $\widetilde{A} = 1$ and convergence constant $\beta = 1$.

n	l	B	C	Shifted $1/N$ E[7]	Moment E[6]	AIM E[14]	Present Work E
0	0	1	1	0.60025	0.59377	0.59365	0.59377

TABLE IV: Comparison of our results with the shifted $1/N$ expansion, moment and perturbative asymptotic iteration methods for the energy eigenvalues, where $A = 0$, $B = 1$, $C = 1$, $\kappa = 2$, m=1 and $\hbar = 1$.

C	E [6]	E_{AIM}
0.1	-0.296088	-0.29608776
0.5	0.1796683	0.17966848
1.0	0.5937711	0.59377126
2.0	1.2237050	1.22370510
5.0	2.5617326	2.56173268
10.0	4.1501236	4.15012364
20.0	6.4799505	6.47995056
50.0	11.2654474	11.26544748
100.0	16.8052478	16.80524784
1000.0	59.3754689	59.37546904
2000.0	85.7348038	85.73480386
5000.0	138.5571975	138.55719764

TABLE V: Comparison of our results with the ground state energy eigenvalues of the hydrogen atom in the $V(r) = Cr^2$ potential, where $A = 0$, $B = 1$, $\kappa = 2$, m=1 and $\hbar = 1$.